

Permutations containing and avoiding certain patterns

Toufik Mansour
Department of Mathematics
Haifa University

February 1, 2008

Abstract

Let $T_k^m = \{\sigma \in S_k \mid \sigma_1 = m\}$. We prove that the number of permutations which avoid all patterns in T_k^m equals $(k-2)!(k-1)^{n+1-k}$ for $k \leq n$. We then prove that for any $\tau \in T_k^1$ (or any $\tau \in T_k^k$), the number of permutations which avoid all patterns in T_k^1 (or in T_k^k) except for τ and contain τ exactly once equals $(n+1-k)(k-1)^{n-k}$ for $k \leq n$. Finally, for any $\tau \in T_k^m$, $2 \leq m \leq k-1$, this number equals $(k-1)^{n-k}$ for $k \leq n$. These results generalize recent results due to Robertson concerning permutations avoiding 123-pattern and containing 132-pattern exactly once.

1 Introduction

In 1990, Herb Wilf asked the following: How many permutations, of length n , avoid a given pattern τ ? By pattern-avoiding we mean the following: A permutation $\alpha \in S_n$ avoids a permutation $\tau \in S_m$ if there is no $1 \leq i_1 < \dots < i_m \leq n$ such that $(\alpha_{i_1}, \dots, \alpha_{i_m})$ is order-isomorphic to $\tau = (\tau_1, \dots, \tau_m)$. In this case, we write $\alpha \in S_n(\tau)$ and call the permutation τ a *pattern*. If a permutation $\alpha \in S_n$ avoids all patterns τ in a set T , we say that α avoids T and write $\alpha \in S_n(T)$.

The first case to be examined was the case of permutations avoiding one pattern of length 3. Knuth [2] found that $|S_n(132)| = |S_n(123)| = c_n$ where c_n is the n -th Catalan number given by the formula $c_n = \frac{1}{n+1} \binom{2n}{n}$, and it

is easy to prove that $|S_n(\tau)| = c_n$ for all $\tau \in S_3$. Later then Simion and Schmidt [6] found the cardinalities of $|S_n(T)|$ for all $T \subseteq S_3$.

A permutation $\alpha \in S_n$ contains $\tau \in S_m$ exactly r times if there exist exactly r different sequences $1 \leq i_1^j < \dots < i_m^j \leq n$, $1 \leq j \leq r$, such that $(\alpha_{i_1^j}, \dots, \alpha_{i_m^j})$ is order-isomorphic to $\tau = (\tau_1, \dots, \tau_m)$ for all $j = 1, \dots, r$.

Noonan shows in [3] that the number of permutations containing exactly one 123-pattern is given by a simple formula $\frac{3}{n} \binom{2n}{n+3}$. Bona [1] proves that expression $\binom{2n-3}{n-3}$ enumerates those permutations containing exactly one 132-pattern, and this result was extended by Robertson, Wilf and Zeilberger [5] to calculate the number of 132-avoiding permutations that have a given number of 123-patterns.

Robertson [4] proves that $(n-2)2^{n-3}$ gives the number of permutations containing exactly one 132-pattern and avoiding the 123-pattern; he also shows that the number of permutations containing exactly one 123-pattern and one 132-pattern equals $(n-3)(n-4)2^{n-5}$.

In this note we obtain a generalization of Robertson's result concerning the number 132-avoiding permutations that have one 123-pattern. As a byproduct we get a generalization of the following result due to Simion and Schmidt [6]: $|S_n(123, 132)| = |S_n(213, 231)| = 2^{n-1}$.

Acknowledgements. I am grateful to R. Chapman and to the anonymous referee for their helpful comments. Special thanks to my advisor A. Vainshtein for his help during the preparation of the paper.

2 Permutations avoiding T_k^m

Let $T_k^m = \{\sigma \in S_k \mid \sigma_1 = m\}$ for all $1 \leq m \leq k$, $k \in \mathcal{N}$. For example $T_3^1 = \{123, 132\}$ and $T_4^2 = \{2134, 2143, 2314, 2341, 2413, 2431\}$.

As a preparatory step, we calculate the cardinalities of the sets $S_n(T_k^m)$ for all $1 \leq m \leq k$, $k \in \mathcal{N}$.

Theorem 2.1

$$|S_n(T_k^m)| = (k-2)! \cdot (k-1)^{n+2-k},$$

for all $m, k, n \in \mathcal{N}$ such that $2 < k \leq n$, $1 \leq m \leq k$.

Proof Let $G_n = S_n(T_k^m)$, and define the family of functions $f_h : S_n \rightarrow S_{n+1}$ by:

$$[f_h(\beta)]_i = \begin{cases} h, & \text{when } i = 1 \\ \beta_{i-1}, & \text{when } \beta_{i-1} < h \\ \beta_{i-1} + 1, & \text{when } \beta_{i-1} \geq h \end{cases}$$

for every $i = 1, \dots, n+1$, $\beta \in S_n$ and $h = 1, \dots, n+1$.

From this we see that:

if $\sigma \in G_n$ then $f_{n+1}(\sigma), f_n(\sigma), \dots, f_{n+m-k+2}(\sigma), f_1(\sigma), \dots, f_{m-1}(\sigma) \in G_{n+1}$
so $(k-1) \cdot |G_n| \leq |G_{n+1}|$ where $n \geq k$.

Assume that $(k-1) \cdot |G_n| < |G_{n+1}|$. Then there exists a permutation $\alpha \in G_{n+1}$ such that $m \leq \alpha_1 \leq n+m-k+1$, so there exist $k-1$ positions $1 < i_1 < \dots < i_{k-1} \leq n+1$ such that the subsequence $\alpha_1, \alpha_{i_1}, \dots, \alpha_{i_{k-1}}$ is order-isomorphic to one of the patterns in T_k^m , which contradicts the definition of G_{n+1} . So $(k-1) \cdot |G_n| = |G_{n+1}|$ for $n \geq k$. Besides $|G_k| = (k-1)(k-1)!$ (from the definition of this set), hence $|G_n| = (k-2)! \cdot (k-1)^{n+2-k}$ for $n \geq k$. ■

Example 2.1 As a corollary we get the result of Simion and Schmidt [6]: $|S_n(T_3^1)| = |S_n(T_3^2)| = 2^{n-1}$ for all $n \in \mathcal{N}$. Other examples: $|S_n(T_4^1)| = 2 \cdot 3^{n-2}$ and $|S_n(T_5^2)| = 3 \cdot 2^{2n-5}$.

Corollary 2.1

$$|S_n(T_k^a \cup T_k^{a+1} \cup \dots \cup T_k^b)| = (k-1)!(k+a-b-1)^{n+1-k},$$

where $1 \leq a \leq b \leq k$.

Proof Let $G_n = S_n(T_k^a \cup T_k^{a+1} \cup \dots \cup T_k^b)$. From Theorem 2.1 we get that $\alpha \in G_n$ if and only if

$$f_1(\alpha), \dots, f_{a-1}(\alpha), f_{n+b-(k-2)}(\alpha), \dots, f_{n+1}(\alpha) \in G_{n+1}.$$

So $|G_{n+1}| = (k+a-b-1)|G_n|$. Besides $|G_k| = (k+a-b+1)(k-1)!$, hence the theorem holds. ■

For a general family $T_k^{i_1}, \dots, T_k^{i_d}$ the situation is more complicated. However, the following recurrence can be proved.

Corollary 2.2 For all $n \geq 2k+1$,

$$|S_n(T_k^{i_1} \cup \dots \cup T_k^{i_d})| = (k+i_1-i_d-1)|S_{n-1}(T_k^{i_1} \cup \dots \cup T_k^{i_d})|,$$

where $1 \leq i_1 < i_2 < \dots < i_d \leq k$.

3 Permutations avoiding $T_k^1 \setminus \{\tau\}$ and containing τ exactly once

Let $b_1 < \dots < b_n$; we denote by $S_{\{b_1, \dots, b_n\}}$ the set of all permutation of the numbers b_1, \dots, b_n ; for example, $S_{\{1, \dots, n\}}$ is just S_n . As above we denote by $S_{\{b_1, \dots, b_n\}}(T)$ the set of all permutations in $S_{\{b_1, \dots, b_n\}}$ avoiding all the permutations in T .

Proposition 3.1 Let $T \subseteq S_{\{c_1, \dots, c_k\}}$. Then there exists $R \subseteq S_k$ such that $|S_n(R)| = |S_{\{c_1, \dots, c_n\}}(T)|$.

Proof We define a function $f : S_{\{c_1, \dots, c_k\}} \rightarrow S_k$ by

$$f((c_{i_1}, c_{i_2}, \dots, c_{i_k})) = (i_1, i_2, \dots, i_k),$$

then evidently $|S_{\{c_1, \dots, c_n\}}(\tau)| = |S_n(f(\tau))|$ for all $\tau \in S_{\{c_1, \dots, c_k\}}$.

Let $T = \{\tau_1, \dots, \tau_l\}$ and $R = \{f(\tau_1), \dots, f(\tau_l)\}$. So

$$S_{\{c_1, \dots, c_n\}}(T) = \bigcap_{i=1}^l S_{\{c_1, \dots, c_n\}}(\tau_i),$$

hence by the isomorphism f we have that $|S_{\{c_1, \dots, c_n\}}(T)| = |S_n(R)|$. ■

Defintion 3.1 Let $M_\tau^{k,m} = T_k^m \setminus \{\tau\}$, for $\tau \in T_k^m$. We denote by $S_n(T_k^m; \tau)$ the set of all permutations in S_n that avoid $M_\tau^{k,m}$ and contain τ exactly once.

Theorem 3.1

$$|S_n(T_k^1; \tau)| = (n+1-k) \cdot (k-1)^{n-k},$$

for all $k \leq n$, $\tau \in T_k^1$.

Proof Let $\alpha \in S_n(T_k^1; \tau)$, and let us consider the possible values of α_1 :

1. $\alpha_1 \geq n-k+2$. Evidently $\alpha \in S_n(T_k^1; \tau)$ if and only if $\alpha \in S_{\{1, \dots, n\} \setminus \{\alpha_1\}}(T_k^1; \tau)$.
2. $\alpha_1 \leq n-k$. Then there exist $1 < i_1 < \dots < i_k \leq n$ such that $(\alpha_1, \alpha_{i_1}, \dots, \alpha_{i_k})$ is a permutation of the numbers $n, \dots, n-k+1, \alpha_1$. For any choice of $k-1$ positions out of i_1, \dots, i_k , the corresponding permutations preceeded by α_1 is order-isomorphic to some permutation in T_k^1 . Since α avoids $M_\tau^{k,1}$, it is, in fact, order-isomorphic to τ . We thus get at least k occurrences of τ in α , a contradiction.

3. $\alpha_1 = n - k + 1$. Then there exist $1 < i_1 < \dots < i_{k-1} \leq n$ such that $\eta = (\alpha_1, \alpha_{i_1}, \dots, \alpha_{i_{k-1}})$ is a permutation of the numbers $n, \dots, n - k + 1$. As above, we immediately get that η is order-isomorphic to τ . We denote by A_n the set of all permutations in $S_n(T_k^1; \tau)$ such that $\alpha_1 = n - k + 1$, and define the family of functions $f_h : A_n \rightarrow S_{n+1}$ by:

$$[f_h(\beta)]_i = \begin{cases} 1, & \text{when } i = h \\ \beta_i + 1, & \text{when } i < h \\ \beta_{i-1} + 1, & \text{when } i > h \end{cases},$$

for every $i = 1, \dots, n + 1$, $\beta \in A_n$ and $h = 1, \dots, n + 1$. It is easy to see that for all $\beta \in A_n$,

$$f_{n+1}(\beta), \dots, f_{n-k+3}(\beta) \in A_{n+1},$$

hence $(k - 1)|A_n| \leq |A_{n+1}|$.

Now we define another function $g : A_{n+1} \rightarrow S_n$ by:

$$[g(\beta)]_i = \begin{cases} \beta_i - 1, & \text{when } i < h \\ \beta_{i+1} - 1, & \text{when } i + 1 > h \end{cases},$$

where $\beta_h = 1$, $i = 1, \dots, n$, $\beta \in A_{n+1}$.

Observe that $h \geq n - k + 3$, since otherwise already $(\beta_h, \beta_{h+1}, \dots, \beta_{n+1})$ contains a pattern from T_k^1 , a contradiction. It is easy to see that $g(\beta) \in A_n$ for all $\beta \in A_{n+1}$, hence $|A_{n+1}| \leq (k - 1)|A_n|$. So finally, $|A_{n+1}| = (k - 1)|A_n|$ and $|A_n| = (k - 1)^{n-k}$, since $|A_k| = 1$.

Since the above cases 1, 2, 3 are disjoint, and by Theorem 2.1 and Proposition 3.1 we obtain

$$|S_n(T_k^1; \tau)| = (k - 1)|S_{n-1}(T_k^1; \tau)| + (k - 1)^{n-k},$$

hence

$$|S_n(T_k^1; \tau)| = (n + 1 - k) \cdot (k - 1)^{n-k},$$

for all $k \leq n$, $\tau \in T_k^1$. ■

Example 3.1 $|S_n(123; 132)| = |S_n(132; 123)| = (n - 2)2^{n-3}$, which is the result of Robertson in [4].

Corollary 3.1

$$|S_n(T_k^k; \tau)| = (n + 1 - k) \cdot (k - 1)^{n-k},$$

for all $k \leq n$, $\tau \in T_k^k$.

Proof Let β be a permutation complement to τ . By the natural bijection between the set $S_n(T_k^1; \beta)$ and the set $S_n(T_k^k, \tau)$ for all $\tau \in T_k^k$ we have that $S_n(T_k^k; \tau)$ have the same cardinality as $S_n(T_k^1; \beta)$, which is $(n + 1 - k) \cdot (k - 1)^{n-k}$ by Theorem 3.1. ■

4 Permutations avoiding $T_k^m \setminus \{\tau\}$ and containing τ exactly once, $2 \leq m \leq k - 1$

Now we calculate the cardinalities of the sets $S_n(T_k^m; \tau)$ where $2 \leq m \leq k - 1$, $\tau \in T_k^m$.

Theorem 4.1

$$|S_n(T_k^m; \tau)| = (k - 1)^{n-k},$$

for all $2 \leq m < k \leq n$, $\tau \in T_k^m$.

Proof Let $G_n = S_n(T_k^m; \tau)$, $\alpha \in G_n$, and let us consider the possible values of α_1 :

1. Let $\alpha_1 \leq m - 1$. Evidently $\alpha \in G_n$ if and only if $(\alpha_2, \dots, \alpha_n) \in S_{\{1, \dots, n\} \setminus \{\alpha_1\}}(T_k^m; \tau)$.
2. $\alpha_1 \geq n - k + m + 1$. Evidently $\alpha \in G_n$ if and only if $(\alpha_2, \dots, \alpha_n) \in S_{\{1, \dots, n\} \setminus \{\alpha_1\}}(T_k^m; \tau)$.
3. $m \leq \alpha_1 \leq n - k + m$. By definition we have that $|G_k| = 1$, so let $n \geq k + 1$. If $m + 1 \leq \alpha_1$ then α contains at least $m \geq 2$ occurrences of a pattern from T_k^m , and If $\alpha_1 \leq n - k + m - 1$ then α contains at least $k - m + 1 \geq 2$ occurrences of a pattern from T_k^m , a contradiction.

Since the above cases 1, 2, 3 are disjoint and by Proposition 3.1 we obtain $|G_n| = (k - 1)|G_{n-1}|$ for all $k \leq n$. Besides $|G_k| = 1$, hence $|S_n(T_k^m; \tau)| = (k - 1)^{n-k}$. ■

Example 4.1 $|S_n(213; 231)| = |S_n(231; 213)| = 2^{n-3}$.

References

- [1] M. BONA, Permutations with one or two 132-subsequences. *Discrete Mathematics*, 181, 1998, 267-274.
- [2] D.E. KNUTH, *The Art of Computer Programming*, 2nd ed. Addison Wesley, Reading, MA, 1973.
- [3] J. NOONAN, The number of permutations containing exactly one increasing subsequence of length 3. *Discrete Mathematics*, 152 (1996), 307-313.
- [4] A. ROBERTSON, Permutations containing and avoiding 123 and 132 patterns, *Discrete Math. Theoret. Computer Science* 3 (1999), 151-154.
- [5] A. ROBERTSON, H.S. WILF and D. ZEILBERGER, Patterns and Fractions, *Electronic Journal of Combinatorics* 6 (1999), #R38.
- [6] R. SIMION AND F.W. SCHMIDT, Restricted permutations, *European Journal of Combinatorics* 6 (1985) 383-406.